Acetyl-HIST1H4A (K91) Antibody



PACO59617

Reactivity:

Source:

Product Information

Size: Protein Background:

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA

Human replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

nucleosome remodeling.

Rabbit Gene ID:

Isotype: HIST1H4A

lgG Uniprot

Applications: P62805

ELISA, WB, IF Synonyms:

Recommended dilutions: Histone H4, HIST1H4A; HIST1H4B; HIST1H4C; HIST1H4E; HIST1H4F;

HIST1H4H; HIST1H4I; HIST1H4I; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; HIST4H4, HIST2H4B; HIST4H4, HIST2H4B; HIST3H4H, HIST2H4B; HIST3H4H, HIST2H4B; HIST3H4H, HIST2H4B; HIST3H4H, HIST2H4B; HIST2H4B

H4/M H4FM; H4/E H4FE; H4/D H4FD; H4/K H4FK; H4/N H4F2 H4FN HIST2H4; H4/O

H4FO;


Immunogen:

Peptide sequence around site of Acetyl-Lys (91) derived from Human Histone H4.


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Detected samples: Hela whole cell lysate, HepG2 whole cell lysate, 293 whole cell lysate, A549 whole cell lysate; Untreated (-) or treated (+) with 30mM sodium butyrate for 4h. All lanes: HIST1H4A antibody at 1:1000. Secondary. Goat polyclonal to rabbit IgG at 1/40000 dilution. Predicted band size: 12 kDa. Observed band size: 12 kDa.



Immunofluorescence staining of Hela cells (treated with 30mM sodium butyrate for 4h) with PACO59617 at 1:5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).