

# Crotonyl-HIST1H2BC (K23) Antibody



PACO59625

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IF, IP, ChIP

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:100-1:1000,  
IF:1:20-1:200, IP:1:200-1:2000

**Protein Background:**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

**Gene ID:**

HIST1H2BC

**Uniprot**

P62807

**Synonyms:**

Histone H2B type 1-C/E/F/G/I (Histone H2B.1 A) (Histone H2B. a) (H2B/a) (Histone H2B. g) (H2B/g) (Histone H2B. h) (H2B/h) (Histone H2B. k) (H2B/k) (Histone H2B. l) (H2B/l), HIST1H2BC; HIST1H2BE; HIST1H2BF; HIST1H2BG; HIST1H2BI, H2BFL; H2BFH; H2BFG; H2BFA; H2BFK


**Immunogen:**

Peptide sequence around site of Crotonyl-Lys (23) derived from Human Histone H2B type 1-C/E/F/G/I.


**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


## Product Images



Western Blot. Positive WB detected in: HeLa whole cell lysate, HL60 whole cell lysate, MCF-7 whole cell lysate, K562 whole cell lysate. All lanes: HIST1H2BC antibody at 1.8 $\mu$ g/ml. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 14 kDa. Observed band size: 14 kDa.



Immunoprecipitating HIST1H2BC in K562 whole cell lysate. Lane 1: Rabbit control IgG instead of PACO59625 in K562 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000). Lane 2: PACO59625 (3 $\mu$ g) + K562 whole cell lysate (500 $\mu$ g). Lane 3: K562 whole cell lysate (20 $\mu$ g).



Immunofluorescence staining of HeLa cells (treated with 30mM sodium crotonylate for 4h) with PACO59625 at 1:37.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).