## Crotonyl-HIST1H2BC (K11) Antibody


## PACO59646



Product Information	
Size:	Protein Background:
50ul	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Reactivity:	
Human	
Source:	
Rabbit	Gene ID:
lsotype:	HIST1H2BC
lgG	Uniprot
Applications:	P62807
ELISA, WB, IF	Synonyms:
Recommended dilutions:	Histone H2B type 1-C/E/F/G/I (Histone H2B.1 A) (Histone H2B. a) (H2B/a) (Histone H2B. g) (H2B/g) (Histone H2B. h) (H2B/h) (Histone H2B. k) (H2B/k) (Histone H2B. l) (H2B/l), HIST1H2BC; HIST1H2BE; HIST1H2BF; HIST1H2BG; HIST1H2BI, H2BFL; H2BFH; H2BFG; H2BFA; H2BFK
ELISA:1:2000-1:10000, WB:1:200-1:2000, IF:1:10-1:100	
	Immunogen:
	Peptide sequence around site of Crotonyl-Lys (11) derived from Human Histone H2B type 1-C/E/F/G/I.
	Storage:

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



Western Blot. Positive WB detected in: Hela whole cell lysate, K562 whole cell lysate, HepG2 whole cell lysate, 293 whole cell lysate (all treated with 30mM sodium crotonylate for 4h). All lanes: HIST1H2BC antibody at 0.36µg/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 14 kDa. Observed band size: 14 kDa.

Immunofluorescence staining of Hela cells (treated with 30mM sodium crotonylate for 4h) with PACO59646 at 1:20, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).