APMAP Antibody



PACO60400

Reactivity:

Product Information

Size: Protein Background:

50ug Exhibits strong arylesterase activity with beta-naphthyl acetate and phenyl acetate. May

play a role in adipocyte differentiation.

Gene ID:

Human, Rat

APMAP

Source:

Uniprot Rabbit

Q9HDC9 **Isotype:**

lgG Synonyms:

Adipocyte plasma membrane-associated protein (Protein BSCv), APMAP, C20orf3

Applications:


ELISA, WB, IHC, IF

Recombinant Human Adipocyte plasma membrane-associated protein (134-240AA).


Recommended dilutions:
Storage:


ELISA:1:2000-1:10000, WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200 Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Positive WB detected in: HepG2 whole cell lysate, Rat liver tissue. All lanes: APMAP antibody at 3.1µg/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 47, 33 kDa. Observed band size: 47 kDa.





IHC image of PACO60400 diluted at 1:400 and staining in paraffinembedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO60400 diluted at 1:400 and staining in paraffinembedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.