

Product Information**Size:**

50ul

Protein Background:

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Reactivity:

Human

Source:

Rabbit

Gene ID:

HIST1H2BC

Isotype:

IgG

Uniprot**Applications:**

ELISA, WB, ICC

Synonyms:**Recommended dilutions:**ELISA:1:2000-1:10000, WB:1:100-1:1000,
ICC:1:20-1:200

Histone H2B type 1-C/E/F/G/I (Histone H2B.1 A) (Histone H2B. a) (H2B/a) (Histone H2B. g) (H2B/g) (Histone H2B. h) (H2B/h) (Histone H2B. k) (H2B/k) (Histone H2B. l) (H2B/l), HIST1H2BC; HIST1H2BE; HIST1H2BF; HIST1H2BG; HIST1H2BI, H2BFL; H2BFH; H2BFG; H2BFA; H2BFK


Immunogen:

Peptide sequence around site of β -hydroxybutyryl-Lys (20) derived from Human Histone H2B type 1-C/E/F/G/I.

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Detected samples: 293 whole cell lysate, A549 whole cell lysate; Untreated (-) or treated (+) with 50mM Sodium 3-hydroxybutyrate for 72h. All lanes: HIST1H2BC antibody at 1:100. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 14 kDa. Observed band size: 14 kDa.

