

HIST1H1D (Ab-85) Antibody



PACO60611

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, CHIP

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:10-1:100

Protein Background:

Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation.

Gene ID:

HIST1H1D

Uniprot

P16402

Synonyms:

Histone H1.3 (Histone H1c) (Histone H1s-2), HIST1H1D, H1F3


Immunogen:

Peptide sequence around site of Lys (85) derived from Human Histone H1.3.


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



IHC image of PACO60611 diluted at 1:20 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Chromatin Immunoprecipitation Hela (4×10^6

) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with $5 \mu\text{g}$ anti-HIST1H1D (PACO60611) or a control normal rabbit IgG. The resulting CHIP DNA was quantified using real-time PCR with primers against the beta -Globin promoter.