
CD164 Antibody

PACO60885



Product Information	
Size:	Protein Background:
50ug	Sialomucin that may play a key role in hematopoiesis by facilitating the adhesion of CD34(+) cells to the stroma and by negatively regulating CD34(+)CD38(lo/-) cell proliferation. Modulates the migration of umbilical cord blood CD133+ cells and this is mediated through the CXCL12/CXCR4 axis. May play an important role in prostate cancer metastasis and the infiltration of bone marrow by cancer cells. Promotes myogenesis by enhancing CXCR4-dependent cell motility. Positively regulates myoblast fusion into myotubes.
Reactivity:	
Human	
Source:	
Rabbit	
lsotype:	Gene ID: CD164 Uniprot Q04900
lgG	
Applications:	
ELISA, IHC, IF	
Recommended dilutions:	Synonyms:
ELISA:1:2000-1:10000, IHC:1:200-1:500, IF:1:50-1:200	Sialomucin core protein 24 (MUC-24) (Endolyn) (Multi-glycosylated core protein 24) (MGC-24) (MGC-24v) (CD antigen CD164), CD164
	Immunogen:
	Recombinant Human Sialomucin core protein 24 protein (24-162AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO60885 diluted at 1:200 and staining in paraffinembedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of A549 cells with PACO60885 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).