

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IF

Recommended dilutions:

ELISA:1:2000-1:10000, IF:1:50-1:200

Protein Background:

Acts as a sulfur carrier required for molybdopterin biosynthesis. Component of the molybdopterin synthase complex that catalyzes the conversion of precursor Z into molybdopterin by mediating the incorporation of 2 sulfur atoms into precursor Z to generate a dithiolene group. In the complex, serves as sulfur donor by being thiocarboxylated (-COSH) at its C-terminus by MOCS3. After interaction with MOCS2B, the sulfur is then transferred to precursor Z to form molybdopterin.

Gene ID:

MOCS2

Uniprot

O96033

Synonyms:

Molybdopterin synthase sulfur carrier subunit (MOCO1-A) (Molybdenum cofactor synthesis protein 2 small subunit) (Molybdenum cofactor synthesis protein 2A) (MOCS2A) (Molybdopterin-synthase small subunit) (Sulfur carrier protein MOCS2A), MOCS2, MOCO1


Immunogen:

Recombinant Human Molybdopterin synthase sulfur carrier subunit protein (1-88AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Immunofluorescence staining of HepG2 cells with PACO60945 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).