ACHE Antibody



PACO60949

Reactivity:

Product Information

Recommended dilutions:

Size: Protein Background:

50ug Terminates signal transduction at the neuromuscular junction by rapid hydrolysis of the

acetylcholine released into the synaptic cleft. Role in neuronal apoptosis.

Gene ID: Human

ACHE

Source: Uniprot

Rabbit P22303

Isotype:

lgG Synonyms:

Acetylcholinesterase (AChE) (EC 3.1.1.7), ACHE Applications:


ELISA, IHC, IF

Recombinant Human Acetylcholinesterase protein (53-162AA).


ELISA:1:2000-1:10000, IHC:1:200-1:500,

IF:1:50-1:200 Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



IHC image of PACO60949 diluted at 1:300 and staining in paraffinembedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with PACO60949 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).