

## Product Information

**Size:**

50ul

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC, IF

**Recommended dilutions:**

ELISA:1:2000-1:10000, IHC:1:500-1:1000,  
IF:1:50-1:200

**Protein Background:**

S-adenosyl-L-methionine-dependent methyltransferase that catalyzes four methylations of the modified target histidine residue in translation elongation factor 2 (EF-2), to form an intermediate called diphthine methyl ester. The four successive methylation reactions represent the second step of diphthamide biosynthesis.

**Gene ID:**

DPH5

**Uniprot**

Q9H2P9

**Synonyms:**

Diphthine methyl ester synthase (EC 2.1.1.314) (Diphthamide biosynthesis methyltransferase), DPH5

**Immunogen:**


Recombinant Human Diphthine methyl ester synthase protein (138-235AA).

**Storage:**


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



IHC image of PACO61722 diluted at 1:500 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of SH-SY5Y cells with PACO61722 at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).