
## **AXIN2 Antibody**

## PACO62119



Product Information	
Size:	Protein Background:
50ul	Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate
Reactivity:	the phosphorylation of beta-catenin and APC by GSK3B.
Human	Gene ID:
Source:	AXIN2
Rabbit	Uniprot
	Q9Y2T1
lsotype:	Synonyms:
lgG	
Applications:	Axin-2 (Axin-like protein) (Axil) (Axis inhibition protein 2) (Conductin), AXIN2
ELISA, IHC, IF	Immunogen:
Recommended dilutions:	Peptide sequence from Human Axin-2 protein (56-73AA).
Recommended dilutions:	Storage:
ELISA:1:2000-1:10000, IHC:1:20-1:200, IF:1:50-1:200	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO62119 diluted at 1:100 and staining in paraffinembedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of U251 cells with PACO62119 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO62119 diluted at 1:100 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.