

PACO62123

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:1000-1:5000,  
IHC:1:20-1:200

**Protein Background:**

**Gene ID:**

ADAR

**Uniprot**

P55265

**Synonyms:**

Double-stranded RNA-specific adenosine deaminase (DRADA) (EC 3.5.4.37) (136 kDa double-stranded RNA-binding protein) (p136) (Interferon-inducible protein 4) (IFI-4) (K88DSRBP), ADAR, ADAR1 DSRAD G1P1 IFI4


**Immunogen:**

Recombinant Human Double-stranded RNA-specific adenosine deaminase protein (367-471AA).


**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images



Western Blot. Positive WB detected in: HeLa whole cell lysate, 293T whole cell lysate, Jurkat whole cell lysate, HepG2 whole cell lysate, U87 whole cell lysate, A549 whole cell lysate, LO2 whole cell lysate. All lanes: ADAR antibody at 1:2000. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 137, 134, 132, 141, 104 kDa. Observed band size: 104 kDa.



IHC image of PAC062123 diluted at 1:30 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.