BCL2L10 Antibody



PACO62623

Product Information

Size: Protein Background:

50ul Promotes cell survival. Suppresses apoptosis induced by BAX but not BAK.

Reactivity: Gene ID:

Human BCL2L10

Source: Uniprot

Rabbit Q9HD36

Isotype: Synonyms:

lgG Bcl-2-like protein 10 (Bcl2-L-10) (Anti-apoptotic protein NrH) (Apoptosis regulator Bcl-

B), BCL2L10, BCLB

Storage:

Applications:

Immunogen:

ELISA, IHC, IF

Recommended dilutions:


ELISA:1:2000-1:10000, IHC:1:20-1:200,

IF:1:20-1:200


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Recombinant Human Bcl-2-like protein 10 protein (19-127AA).


Product Images



IHC image of PACO62623 diluted at 1:150 and staining in paraffinembedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of A549 cells with PACO62623 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of PACO62623 diluted at 1:150 and staining in paraffinembedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.