UNC13B Antibody



PACO62723

Rabbit

Isotype:

Product Information

Size: **Protein Background:**

50ul Plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol

second messenger pathway. Is involved in neurotransmitter release by acting in

Reactivity: synaptic vesicle priming prior to vesicle fusion and participates in the activity-

depending refilling of readily releasable vesicle pool (RRP). Essential for synaptic vesicle Human

maturation in a subset of excitatory/glutamatergic but not inhibitory/GABA-mediated

Source: synapses.

lgG Uniprot

O14795 **Applications:**

ELISA, WB, IHC, IF Synonyms:

Protein unc-13 homolog B (Munc13-2) (munc13), UNC13B, UNC13 **Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:1000-1:5000,

IHC:1:200-1:500, IF:1:50-1:200

Immunogen:

Recombinant Human Protein unc-13 homolog B protein (12-215AA).


Storage:

Gene ID:


UNC13B

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


Product Images



IHC image of PACO62723 diluted at 1:300 and staining in paraffinembedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot. Positive WB detected in: U251 whole cell lysate, MCF-7 whole cell lysate. All lanes: UNC13B antibody at 1:2000. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 181, 183 kDa. Observed band size: 181 kDa.



Immunofluorescence staining of Hela cells with PACO62723 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).