FLT3 Antibody, Biotin conjugated

PACO62898



Product Information	
Size:	Protein Background:
50ul	Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and
Reactivity:	regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of
Human	the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1.
Source:	Promotes phosphorylation of FES, FER, PTPN6/SHP, PTPN11/SHP-2, PLCG1, and
Rabbit	STAT5A and/or STAT5B. Activation of wild-type FLT3 causes only marginal activation of STAT5A or STAT5B. Mutations that cause constitutive kinase activity promote cell
lsotype:	proliferation and resistance to apoptosis via the activation of multiple signaling pathways.
lgG	Gene ID:
Applications:	FLT3
ELISA	Uniprot
Recommended dilutions:	P36888
	Synonyms:
	Receptor-type tyrosine-protein kinase FLT3 (EC 2.7.10.1) (FL cytokine receptor) (Fetal liver kinase-2) (FLK-2) (Fms-like tyrosine kinase 3) (FLT-3) (Stem cell tyrosine kinase 1) (STK-1) (CD antigen CD135), FLT3, CD135 FLK2 STK1
	Immunogen:
	- Recombinant Human Receptor-type tyrosine-protein kinase FLT3 protein (727-993AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A