

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200,
IF:1:50-1:200

Protein Background:

autophagosome, cytosol, perinuclear endoplasmic reticulum, plasma membrane, cholesterol binding, sterol transporter activity, bile acid, biosynthetic process, cellular response to cholesterol, positive regulation of proteasomal protein catabolic process, regulation of autophagy.

Gene ID:

OSBPL7

Uniprot

Q9BZF2

Synonyms:

Oxysterol-binding protein-related protein 7 (ORP-7) (OSBP-related protein 7), OSBPL7, ORP7


Immunogen:

Recombinant Human Oxysterol-binding protein-related protein 7 protein (1-249AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



IHC image of PACO63487 diluted at 1:100 and staining in paraffin-embedded human small intestine tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HeLa cells with PACO63487 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).