## PLG Antibody, HRP conjugated



## PACO63696

Reactivity:

Source:

Rabbit

## **Product Information**

Size: Protein Background:

50ul Plasmin dissolves the fibrin of blood clots and acts as a proteolytic factor in a variety of

other processes including embryonic development, tissue remodeling, tumor invasion, and inflammation. In ovulation, weakens the walls of the Graafian follicle. It activates the

Human urokinase-type plasminogen activator, collagenases and several complement

zymogens, such as C1 and C5. Cleavage of fibronectin and laminin leads to cell detachment and apoptosis. Also cleaves fibrin, thrombospondin and von Willebrand

factor. Its role in tissue remodeling and tumor invasion may be modulated by CSPG4.

Binds to cells.

Isotype: Gene ID:

lgG PLG

Applications: Uniprot

ELISA P00747

Recommended dilutions: Synonyms:

Plasminogen (EC 3.4.21.7) [Cleaved into: Plasmin heavy chain A; Activation peptide;

Angiostatin; Plasmin heavy chain A, short form; Plasmin light chain B], PLG

Immunogen:

Recombinant Human Plasminogen protein (103-343AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A