SNF8 Antibody, FITC conjugated



PACO63869

Human

Product Information

Recommended dilutions:

Size: Protein Background:

50ul Component of the endosomal sorting complex required for transport II (ESCRT-II), which is required for multivesicular body (MVB) formation and sorting of endosomal Reactivity:

cargo proteins into MVBs. The MVB pathway mediates delivery of transmembrane proteins into the lumen of the lysosome for degradation. The ESCRT-II complex is

probably involved in the recruitment of the ESCRT-III complex. The ESCRT-II complex

Source: may also play a role in transcription regulation by participating in derepression of

Rabbit transcription by RNA polymerase II, possibly via its interaction with ELL. Required for degradation of both endocytosed EGF and EGFR, but not for the EGFR ligand-mediated

Isotype: internalization. It is also required for the degradation of CXCR4. Required for the

exosomal release of SDCBP, CD63 and syndecan.

Q96H20

lgG **Gene ID:**

Applications: SNF8

ELISA Uniprot

Synonyms:

Vacuolar-sorting protein SNF8 (ELL-associated protein of 30 kDa) (ESCRT-II complex subunit VPS22) (hVps22), SNF8, EAP30

Immunogen:

Recombinant Human Vacuolar-sorting protein SNF8 protein (1-258AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

| Product | Images |
|---------|---------------|
|---------|---------------|

N/A N/A