Phospho-EGFR (Y1068) Recombinant Antibody



RACO0095

Source:

Product Information

Size: **Protein Background:**

50ul Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses.

Reactivity: Known ligands include EGF, TGFA/TGF-alpha, amphiregulin, epigen/EPGN,

BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding Human

> triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2

which in turn activates complex downstream signaling cascades. Human

Gene ID: Isotype:

EGFR Rabbit IgG

Uniprot **Applications:**

P00533 ELISA, WB

Synonyms: **Recommended dilutions:**

Epidermal growth factor receptor, Proto-oncogene c-ErbB-1, Receptor tyrosine-protein WB:1:500-1:5000 kinase erbB-1, EGFR, ERBB, ERBB1, HER1


A synthesized peptide derived from human Phospho-EGFR (Y1068).

Storage:

Immunogen:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Product Images



Western Blot

Positive WB detected in(A549 whole cell lysate) HepG2 whole cell lysate)

(treated with EGF or Pervanadate)

All lanes: Phospho-EGFR antibody at 1.28µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 175 KDa Observed band size: 175 KDa