GRIA2/GRIA3 Recombinant Antibody

RACO0164



Product Information	
Size:	Protein Background:
50ul	Receptor for glutamate that functions as ligand-gated ion channel in the central
Reactivity:	nervous system and plays an important role in excitatory synaptic transmission. L- glutamate acts as an excitatory neurotransmitter at many synapses in the central
Human	nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby
Source:	converts the chemical signal to an electrical impulse. The receptor then desensitizes
Human	rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization
lsotype:	which is characterized by a delayed accumulation of current flux upon continued application of glutamate. Through complex formation with NSG1, GRIP1 and STX12
Rabbit IgG	controls the intracellular fate of AMPAR and the endosomal sorting of the GRIA2
Applications:	subunit toward recycling and membrane targeting (By similarity).
ELISA, IF	Gene ID:
Recommended dilutions:	GRIA2/GRIA3
	Uniprot
IF:1:20-1:200	P42262/P42263
	Synonyms:
	Glutamate receptor 2, GluR-2, AMPA-selective glutamate receptor 2, GluR-B, GluR-K2, Glutamate receptor ionotropic, AMPA 2, GluA2, GRIA2, GLUR2, Glutamate receptor 3, GluR-3, AMPA-selective glutamate receptor 3, GluR-C, GluR-K3, Glutamate receptor ionotropic, AMPA 3, GluA3, GRIA3, GLUR3, GLURC

Immunogen:

A synthesized peptide derived from human GRIA2/GRIA3.

Storage:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.



Immunofluorescence staining of MCF-7 cells with RACO0164 at 1:56, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG (H+L).