MMP9 Recombinant Antibody



RACO0183

Product Information

IHC:1:50-1:200

Size: Protein Background:

50ul May play an essential role in local proteolysis of the extracellular matrix and in

Reactivity: leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at

a Gly-|-Leu bond. Cleaves type IV and type V collagen into large C-terminal three

Human quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin

but not laminin or Pz-peptide.

Source: Gene ID:

Human MMP9

Isotype: Uniprot

Rabbit IgG P14780

Applications: Synonyms:

ELISA, IHC, FC
Matrix metalloproteinase-9, MMP-9, 92 kDa gelatinase, 92 kDa type IV collagenase,

Recommended dilutions: Gelatinase B, GELB, 67 kDa matrix metalloproteinase-9, 82 kDa matrix

metalloproteinase-9, MMP9, CLG4B

Immunogen:

A synthesized peptide derived from human MMP9.


Storage:


Rabbit lgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Copyright © 2021 Assay Genie

info@assaygenie.com

Product Images





Overlay histogram showing Jurkat cells stained with RACO0183 (red line) at for 4h). The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4°C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4°C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.

IHC image of RACO0183 diluted at 1:235 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.