

# TNFRSF12A Recombinant Antibody



RACO0266

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human

**Source:**

Homo sapiens (Human)

**Isotype:**

Rabbit IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

WB:1:500-1:5000, IHC:1:50-1:200

**Protein Background:**

Receptor for TNFSF12/TWEAK. Weak inducer of apoptosis in some cell types. Promotes angiogenesis and the proliferation of endothelial cells. May modulate cellular adhesion to matrix proteins.

**Gene ID:**

TNFRSF12A

**Uniprot**

Q9NP84

**Synonyms:**

Tumor necrosis factor receptor superfamily member 12A (Fibroblast growth factor-inducible immediate-early response protein 14) (FGF-inducible 14) (Tweak-receptor) (TweakR) (CD antigen CD266), TNFRSF12A, FN14

**Immunogen:**


A synthesized peptide derived from human TWEAKR.

**Storage:**


Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

## Product Images

---



IHC image of RACO0266 diluted at 1:100 and staining in paraffin-embedded human placenta tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



### Western Blot

Positive WB detected in( U251 whole cell lysate) MCF-7 whole cell lysate) U87 whole cell lysate) All lanes: TNFRSF12A antibody at 1:2000 Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 14, 11 kD

Observed band size: 14 kDa