
SOX10 Recombinant Antibody

RAC00271



Product Information	
Size:	Protein Background:
50ul	Transcription factor that seems to function synergistically with the POU domain protein
Reactivity:	TST-1/OCT6/SCIP. Could confer cell specificity to the function of other transcription factors in developing and mature glia (By similarity).
Human	Gene ID:
Source:	SOX10
Homo sapiens (Human)	Uniprot
lsotype:	P56693
Rabbit IgG	Synonyms:
Applications:	Transcription factor SOX-10, SOX10
ELISA, WB, IHC, IF	Immunogen:
Recommended dilutions:	A synthesized peptide derived from human SOX10.
WB:1:500-1:5000, IHC:1:50-1:200, IF:1:20- 1:200	Storage:
	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.



IHC image of RACO0271 diluted at 1:100 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

IHC image of RACO0271 diluted at 1:100 and staining in paraffinembedded human melanoma cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Western Blot

Positive WB detected in(Hela whole cell lysate) MCF-7 whole cell lysate) A549 whole cell lysate) HepG2 whole cell lysate) All lanes: SOX10 antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1:50000 dilution Predicted band size: 50, 32 kDa

Observed band size: 60 kDa