HDAC2 Recombinant Antibody



RACO0279

Product Information

Size: **Protein Background:**

50ul Responsible for the deacetylation of lysine residues on the N-terminal part of the core

histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic Reactivity: repression and plays an important role in transcriptional regulation, cell cycle

progression and developmental events. Histone deacetylases act via the formation of Human

large multiprotein complexes. Forms transcriptional repressor complexes by associating Source: with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with

DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1,

Homo sapiens (Human) PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity.

Isotype: Gene ID:

Rabbit IgG HDAC2

Applications: Uniprot

ELISA, WB, IHC Q92769

Recommended dilutions: Synonyms:

WB:1:500-1:5000, IHC:1:50-1:200 Histone deacetylase 2 (HD2) (EC 3.5.1.98), HDAC2


Immunogen:


A synthesized peptide derived from human HDAC2.

Storage:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Product Images





IHC image of RACO0279 diluted at 1:100 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Western Blot

Positive WB detected in (Hela whole cell lysate) HepG2 whole cell lysate) Jurkat whole cell lysate) MCF-7 whole cell lysate) K562 whole cell lysate) All lanes: HDAC2 antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 56, 52 kDa Observed band size: 60 kDa