

# DHFR Recombinant Antibody



RACO0356

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse, Rat

**Source:**

Homo sapiens (Human)

**Isotype:**

Rabbit IgG

**Applications:**

ELISA, WB, IHC, IP

**Recommended dilutions:**

WB:1:500-1:5000, IHC:1:50-1:200, IP:1:200-1:1000

**Protein Background:**

Key enzyme in folate metabolism. Contributes to the de novo mitochondrial thymidylate biosynthesis pathway. Catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis. Binds its own mRNA and that of DHFR2.

**Gene ID:**

DHFR

**Uniprot**

P00374

**Synonyms:**

Dihydrofolate reductase (EC 1.5.1.3), DHFR




**Immunogen:**

A synthesized peptide derived from human DHFR.

**Storage:**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

## Product Images



IHC image of RACO0356 diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

IHC image of RACO0356 diluted at 1:100 and staining in paraffin-embedded human liver cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

### Western Blot

Positive WB detected in( Hela whole cell lysate) Jurkat whole cell lysate) K562 whole cell lysate) Raji whole cell lysate) 293T whole cell lysate) Rat heart tissue, Mouse kidney tissue, Mouse liver tissue

All lanes: DHFR antibody at 1:2000

### Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 22, 16 kDa

Observed band size: 22 kDa