GATA3 Recombinant Antibody



RACO0459

Reactivity:

Product Information

Size: Protein Background:

50ul Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and

delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper

2 (Th2) differentiation process following immune and inflammatory responses.

Human Gene ID:

Source: GATA3

Homo sapiens (Human) Uniprot

Isotype: P23771

Rabbit IgG Synonyms:

Applications: Trans-acting T-cell-specific transcription factor GATA-3 (GATA-binding factor 3), GATA3

ELISA, WB, IHC Immunogen:

Recommended dilutions: A synthesized peptide derived from human GATA3.


WB:1:500-1:5000, IHC:1:50-1:200


Storage:


Rabbit lgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and

50% glycerol.

Product Images







IHC image of RACO0459 diluted at 1:100 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

IHC image of RACO0459 diluted at 1:100 and staining in paraffinembedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Western Blot

Positive WB detected in(RAW264.7 whole cell lysate) All lanes: GATA3

antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 48, 49 kDa Observed band size: 48 kDa