CASP2 Recombinant Antibody



RACO0550

Reactivity:

Product Information

Size: Protein Background:

50ul Involved in the activation cascade of caspases responsible for apoptosis execution.

Might function by either activating some proteins required for cell death or inactivating

proteins necessary for cell survival.

Human Gene ID:

Source: CASP2

Homo sapiens (Human) Uniprot

Isotype: P42575

Rabbit IgG Synonyms:

Applications: Caspase-2 (CASP-2) (EC 3.4.22.55) (Neural precursor cell expressed developmentally

ELISA, WB, IHC down-regulated protein 2) (NEDD-2) (Protease ICH-1) [Cleaved into: Caspase-2 subunit

p18, Caspase-2 subunit p13, Caspase-2 subunit p12], CASP2, ICH1 NEDD2

Recommended dilutions: Immunogen:


WB:1:500-1:5000, IHC:1:50-1:200 A synthesized peptide derived from human Caspase-2.


Storage:


Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and

50% glycerol.

Product Images







IHC image of RACO0550 diluted at 1:100 and staining in paraffinembedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

IHC image of RACO0550 diluted at 1:100 and staining in paraffinembedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Western Blot

Positive WB detected in (K562 whole cell lysate) 293 whole cell lysate)

All lanes: Caspase-2 antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1:50000 dilution

Predicted band size: 51, 35, 11 kDa Observed band size: 51 kDa