ACVR2B Recombinant Antibody

RAC00580



Product Information	
Size:	Protein Background:
50ul	Transmembrane serine/threonine kinase activin type-2 receptor forming an activin
Reactivity:	receptor complex with activin type-1 serine/threonine kinase receptors (ACVR1, ACVR1B or ACVR1c). Transduces the activin signal from the cell surface to the
Human	cytoplasm and is thus regulating many physiological and pathological processes including neuronal differentiation and neuronal survival, hair follicle development and
Source:	cycling, FSH production by the pituitary gland, wound healing, extracellular matrix
Homo sapiens (Human)	production, immunosuppression and carcinogenesis.
lsotype:	Gene ID:
Rabbit IgG	ACVR2B
-	Uniprot
Applications: ELISA, IHC	Q13705
	Synonyms:
Recommended dilutions:	Activin receptor type-2B (EC 2.7.11.30) (Activin receptor type IIB) (ACTR-IIB), ACVR2B
IHC:1:50-1:200	
	Immunogen:
	A synthesized peptide derived from human Activin Receptor Type IIB.
	Storage:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.



IHC image of RACO0580 diluted at 1:100 and staining in paraffinembedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

IHC image of RACO0580 diluted at 1:100 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat antirabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.