

## **Technical Manual**

## **ColorFluor Pyruvate Assay Kit**

**Catalogue Code: BA0146**

**Pack Size: 100 assays**

**Research Use Only**

## DESCRIPTION

**PYRUVATE** is a key intermediate in cellular metabolic pathways. Pyruvate can be converted to carbohydrates via gluconeogenesis, to fatty acids or energy through acetyl-CoA, to the amino acid alanine and to ethanol. Abnormal levels of pyruvate have been linked to liver diseases and metabolic disorders. Simple, direct and automation-ready procedures for measuring pyruvate concentrations find wide applications in research and drug discovery.

The Assay Genie ColorFluor Pyruvate Assay Kit uses a single Working Reagent that combines pyruvate oxidase and hydrogen peroxide determination in one step. The color intensity of the reaction product at 570nm or fluorescence intensity at  $\lambda_{em}/\lambda_{ex} = 585/530\text{nm}$  is directly proportional to pyruvate concentration in the sample.

## KEY FEATURES

**Sensitive and accurate.** Use as little as 10  $\mu\text{L}$  samples. Linear detection range in 96-well plate: 2 to 500  $\mu\text{M}$  (17  $\mu\text{g}/\text{dL}$  to 4.4  $\text{mg}/\text{dL}$ ) pyruvate for colorimetric assays and 0.2 to 50  $\mu\text{M}$  for fluorimetric assays.

**Simple and convenient.** The procedure involves addition of a single working reagent and incubation for 30 min at room temperature, compatible for HTS assays.

**Improved reagent stability.** The optimized formulation has greatly enhanced the reagent and signal stability.

## APPLICATIONS:

**Direct Assays:** pyruvate in biological samples.

**Drug Discovery/Pharmacology:** effects of drugs on pyruvate metabolism.

## KIT CONTENTS

**Enzyme Mix:** 10 mL

**Dye Reagent:** 120  $\mu\text{L}$

**Standard:** 400  $\mu\text{L}$  25 mM Pyruvate

**Storage conditions.** The kit is shipped on dry ice. Store all reagents at  $-20^{\circ}\text{C}$ . Shelf life of six months after receipt.

**Precautions:** reagents are for research use only. Normal precautions for laboratory reagents should be exercised while using the reagents. Please refer to Material Safety Data Sheet for detailed information.

## COLORIMETRIC PROCEDURE

*Note:* SH-group containing reagents (e.g. mercaptoethanol, DTT) may interfere with this assay and should be avoided in sample preparation.

1. Equilibrate all components to room temperature. Prepare a 500  $\mu\text{M}$  Standard Premix by mixing 10  $\mu\text{L}$  of the 25 mM Standard and 490  $\mu\text{L}$   $\text{H}_2\text{O}$ . Dilute Standard in distilled water as follows.

No	Premix + $\text{H}_2\text{O}$	Vol ( $\mu\text{L}$ )	Pyruvate ( $\mu\text{M}$ )
1	100 $\mu\text{L}$ + 0 $\mu\text{L}$	100	500
2	80 $\mu\text{L}$ + 20 $\mu\text{L}$	100	400
3	60 $\mu\text{L}$ + 40 $\mu\text{L}$	100	300
4	40 $\mu\text{L}$ + 60 $\mu\text{L}$	100	200
5	30 $\mu\text{L}$ + 70 $\mu\text{L}$	100	150
6	20 $\mu\text{L}$ + 80 $\mu\text{L}$	100	100
7	10 $\mu\text{L}$ + 90 $\mu\text{L}$	100	50
8	0 $\mu\text{L}$ + 100 $\mu\text{L}$	100	0

Transfer 10  $\mu\text{L}$  standards and 10  $\mu\text{L}$  samples into separate wells of a clear flat-bottom 96-well plate.

2. For each reaction well, mix 94  $\mu\text{L}$  Enzyme Mix and 1  $\mu\text{L}$  Dye Reagent in a clean tube. Transfer 90  $\mu\text{L}$  Working Reagent into each assay well. Tap plate to mix. Freeze unused reagents for future use.

3. Incubate 30 min at room temperature. Read optical density at 570nm (550-585nm).

Note: if the Sample OD is higher than the Standard OD at 500  $\mu\text{M}$ , dilute sample in water and repeat the assay. Multiply result by the dilution factor.

### CALCULATION

Subtract blank OD (water, #8) from the standard OD values and plot the OD against standard concentrations. Determine the slope using linear regression fitting. The pyruvate concentration of Sample is calculated as

$$[\text{Pyruvate}] = \frac{\text{OD}_{\text{SAMPLE}} - \text{OD}_{\text{H}_2\text{O}}}{\text{Slope}} \quad (\mu\text{M})$$

$\text{OD}_{\text{SAMPLE}}$  and  $\text{OD}_{\text{H}_2\text{O}}$  are optical density values of the sample and water.

*Conversions:* 1mM pyruvate equals 8.7 mg/dL or 87 ppm.

### FLUORIMETRIC PROCEDURE

For fluorimetric assays, the linear detection range is 0.2 to 50  $\mu\text{M}$  pyruvate.


- Dilute the Standards prepared in Colorimetric Procedure 1:10 in  $\text{H}_2\text{O}$ .
- Transfer 10  $\mu\text{L}$  standards and 10  $\mu\text{L}$  samples into separate wells of a *black* 96-well plate.
- Add 90  $\mu\text{L}$  Working Reagent (see *Colorimetric Procedure*). Tap plate to mix.
- Incubate 30 min at room temperature and read fluorescence at  $\lambda_{\text{ex}} = 530\text{nm}$  and  $\lambda_{\text{em}} = 585\text{nm}$ .
- If assays in 384-well plate are desired, use 5 $\mu\text{L}$  Standards and 45  $\mu\text{L}$  Working Reagent.

The pyruvate concentration of Sample is calculated as

$$[\text{Pyruvate}] = \frac{F_{\text{SAMPLE}} - F_{\text{H}_2\text{O}}}{\text{Slope}} \quad (\mu\text{M})$$

### MATERIALS REQUIRED, BUT NOT PROVIDED

Pipeting devices, centrifuge tubes, Clear flat-bottom 96-well plates, black 96-well or 384-well plates (e.g. Corning Costar) and plate reader.



### LITERATURE

1. Hansen JL, Freier EF. (1978). Direct assays of lactate, pyruvate, beta-hydroxybutyrate, and acetoacetate with a centrifugal analyzer.

Clin Chem. 24(3):475-9.

2. Sutherland DV, Barns AM, Ross CA. (1995). Trypanosoma evansi: measurement of pyruvate production as an indicator of the drug sensitivity of isolates in vitro. Trop Med Parasitol. 46(2):93-8.

3. Chariot P. et al (1994). Optimal handling of blood samples for routine measurement of lactate and pyruvate. Arch Pathol Lab Med. 118(7):695-7.

## Contact Details

Dublin, Ireland

**Email:** [info@assaygenie.com](mailto:info@assaygenie.com)

**Web:** [www.assaygenie.com](http://www.assaygenie.com)

**Technical Support:** [Techsupport@assaygenie.com](mailto:Techsupport@assaygenie.com)

Copyright © 2017 ReagentBio, All Rights Reserved. All information / detail is correct at time of going to print.