

## Recombinant 2019-nCoV Spike Protein (RBD, His Tag)(Active)

**Catalog No:** RPES0041

**Category:** Recombinant Protein

**Bio-Activity:** Measured by its binding ability in a functional ELISA. Immobilized human ACE2 protein (Fc tag) (Cat# RPES0054) at 2 µg/mL (100 µL/well) can bind Recombinant 2019-nCoV Spike Protein (RBD, His Tag) (Cat#RPES0041), the EC50 of RPES0041 is 35-70 ng/mL.

### Sequence Information

**Species:** Virus

**Sequence:** Arg319-Phe541

**Accession:** YP\_009724390.1

**Tag:** C-His

### Product Information

**Synonyms:** 2019-nCov S protein; 2019-nCoV Spike glycoprotein; 2019-nCoV S glycoprotein

**Source:** Baculovirus-Insect Cells

**Purity:** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per µg as determined by the LAL method.

**Formulation:** Lyophilized from sterile 20mM PB,300mM NaCl, pH 7.0,10% glycerol.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

### Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, Oacetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the


Contact Details | Dublin, Ireland

Email: [hello@assaygenie.com](mailto:hello@assaygenie.com) | Web: [www.assaygenie.com](http://www.assaygenie.com)

Copyright © 2020 Reagent Genie, All Rights Reserved. All information / detail is correct at time of going to print.

membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

**Image**



Contact Details | Dublin, Ireland

Email: [hello@assaygenie.com](mailto:hello@assaygenie.com) | Web: [www.assaygenie.com](http://www.assaygenie.com)

Copyright © 2020 Reagent Genie, All Rights Reserved. All information / detail is correct at time of going to print.