



Recombinant Protein Technical Manual

Recombinant Human EGFR/ErbB1 Protein (His & Avi Tag)

RPE0128

Product Data:

Product SKU: RPE0128

Size: 20µg

Species: Human

Expression host: Human Cells

Uniprot: P00533

Protein Information:

Molecular Mass: 41.5 kDa

AP Molecular Mass: 60-90 kDa

Tag: C-His-Avi

Bio-activity:

Purity: > 95% as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Epidermal growth factor receptor; EGFR; Proto-oncogene c-ErbB; Receptortyrosine-protein kinase erbB; EGFR

Immunogen Information:

Sequence: Leu25-Val30Gly&Asn298-Ser645

Background:

The EGFR subfamily of receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43% - 44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glycoproteins with an extracellular ligand binding domain. The extracellular ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF-alpha, betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.