



Recombinant Protein Technical Manual  
Recombinant Mouse UCHL3/UCH-L3 Protein (His  
Tag)(Active)  
RPES0177

Product Data:

**Product SKU:** RPES0177

**Size:** 20µg

**Species:** Mouse

**Expression host:** E. coli

**Uniprot:** Q9JKB1

Protein Information:

**Molecular Mass:** 27.5 kDa

**AP Molecular Mass:** 30 kDa

**Tag:** N-His

**Bio-activity:** Measured by the hydrolysis of UbiquitinAMC. The specific activity is >14000 pmoles/min/µg.

**Purity:** > 97 % as determined by SDS-PAGE

**Endotoxin:** Please contact us for more information.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from sterile 50mM Tris, 150mM NaCl, 20% glycerol, pH 7.7

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:**

**Synonyms:** UCHL3

## Immunogen Information:

**Sequence:** Glu 2-Ala 230

## Background:

Ubiquitin carboxyl-terminal hydrolase isozyme L3, also known as UCH-L3, Ubiquitin thioesterase L3 and UCHL3, is a ubiquitin-protein hydrolase which belongs to the peptidase C12 family. It is involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. UCHL3 is highly expressed in heart, skeletal muscle, and testis. UCHL1 and UCHL3 are two of the deubiquitinating enzymes expressed in the brain. These phenotypes indicate the importance of UCHL1 and UCHL3 in the regulation of the central nervous system. UCHL3 functions as a de-ubiquitinating enzyme where lack of its hydrolase activity may result in the prominent accumulation of ubiquitinated proteins and subsequent induction of stress responses in skeletal muscle. UCHL3 has also been identified as a tumor-specific antigen in colon cancer.