



Recombinant Protein Technical Manual

Recombinant Rat ICAM/CD54 Protein (His Tag)(Active)
RPES0223

Product Data:

Product SKU: RPES0223

Size: 50µg

Species: Rat

Expression host: HEK293 Cells

Uniprot: Q00238

Protein Information:

Molecular Mass: 52.7 kDa

AP Molecular Mass: 70-75 kDa

Tag: C-His

Bio-activity: Measured by the ability of the immobilized protein to support the adhesion of PMA-stimulated HSB2 human peripheral blood acute lymphoblastic leukemia cells. When cells are added to rat ICAM1 coated plates (12.5 µg/mL, 100 µL/well), approximately > 30% cells will adhere specifically

Purity: > 97 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: CD54;ICAM

Immunogen Information:

Sequence: Met 1-Thr 493

Background:

Intercellular adhesion molecule (ICAM, or CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis. Additionally, ICAM has been implicated in the progression of autoimmune diseases. ICAM is a ligand for LFA(integrin). When activated, leukocytes bind to endothelial cells via ICAM/LFA interaction and then transmigrate into tissues. Presence with heavy glycosylation and other structural characteristics, ICAM possesses binding sites for a number of immune-associated ligands and serves as the binding site for entry of the major group of human Rhinovirus (HRV) into various cell types. ICAM also becomes known for its affinity for Plasmodium falciparum-infected erythrocytes (PFIE), providing more of a role in infectious disease. Previous studies have shown that ICAM is involved in inflammatory reactions and that a defect in ICAM gene inhibits allergic contact hypersensitivity.