



Recombinant Protein Technical Manual  
Recombinant Human PD-L2/CD273 Protein (His Tag)  
RPES0324

### Product Data:

**Product SKU:** RPES0324

**Size:** 10µg

**Species:** Human

**Expression host:** Human Cells

**Uniprot:** Q9BQ51

### Protein Information:

**Molecular Mass:** 23.59 kDa

**AP Molecular Mass:** 34-50 kDa

**Tag:** C-His

**Bio-activity:**

**Purity:** > 95% as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per µg as determined by the LAL method.

**Storage:** Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:**

**Synonyms:** Programmed Cell Death 1 Ligand 2; PD Ligand 2; PD-L2; PDCD1 Ligand 2; Programmed Death Ligand 2; Butyrophilin B7-DC; B7-DC; CD273; PDCD1LG2; B7DC; CD273; PDCD1L2; PDL2

## Immunogen Information:

**Sequence:** Leu20-Pro219

## Background:

Programmed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG family. PDCD1LG2 contains one Ig-like C2-type domain and one Ig-like V-type domain. PDCD1LG2 is highly expressed in the heart, placenta, pancreas, lung and liver; it is weakly expressed in the spleen, lymph nodes, and thymus. PDCD1LG2 is involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.