

Recombinant Protein Technical Manual

Recombinant Mouse CD86/B7-2 Protein (His Tag)(Active) RPES0526

Product Data:

Product SKU: RPES0526 **Size:** 100μg

Species: Mouse Expression host: HEK293 Cells

Uniprot: NP 062261.3

Protein Information:

Molecular Mass: 26.8 kDa

AP Molecular Mass: 40-55 kDa

Tag: C-His

Bio-activity: Measured by its binding ability in a functional ELISA. Immobilized mouse CD86 at

20 μg/ml (100 μl/well) can bind mouse CD28 with a linear ranger of 6.4-800 ng/ml.

Purity: > 92 % as determined by SDS-PAGE

Endotoxin: $< 1.0 \text{ EU per } \mu \text{g}$ of the protein as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: T-lymphocyte activation antigen CD86; Activation B7-2 antigen; Early T-cell

costimulatory molecule 1; ETC; CD86;B7;B7-2;B70;Cd28l2;CLS1;ETC;Ly-

58;Ly58;MB7;MB7-2;TS/A-2

Immunogen Information:

Sequence: Met 1-Glu 245

Background:

CD86, also known as B-lymphocyte activation antigen B7-2 (referred to as B70), is a member of the cell surface immunoglobulin superfamily. B7-2 exists predominantly as a monomer on cell surfaces and interacts with two co-stimulatory receptors CD28 and cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expressed on T cells, and thus induces the signal pathways which regulate T cell activation and tolerance, cytokine production, and the generation of CTL. It is indicated that contacts between B and T helper cells mediated by CD86 encourage signals for the proliferation and IgG secretion of normal B cells and B cell lymphomas. Recent study has revealed that CD86 also promotes the generation of a mature APC repertoire and promotes APC function and survival. CD86 has an important role in chronic hemodialysis, allergic pulmonary inflammation, arthritis, and antiviral responses, and thus is regarded as a promising candidate for immune therapy.