

Recombinant Protein Technical Manual Recombinant Mouse Exostosin-Like 2/EXTL2 Protein (His Tag) RPES0590

Product Data:

Product SKU: RPES0590

Species: Mouse

Size: 10µg

Expression host: Human Cells

Uniprot: Q9ES89

Protein Information:

Molecular Mass:	33.6 kDa
AP Molecular Mass:	35 kDa
Tag:	N-6His
Bio-activity:	
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU per μg as determined by the LAL method.
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris,150mM NaCl,pH8.0.
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	Exostosin-like 2; Extl2; Alpha;4-N-acetylhexosaminyltransferase EXTL2; Alpha- GalNAcT EXTL2; EXT-related protein 2; Glucuronyl-galactosyl-proteoglycan 4- alpha-N-acetylglucosaminyltransferase

Sequence: Asn43-Met330

Background:

Exostosin-like 2 (EXTL2) is a member of the exostosin (EXT)-related family which contains five members: EXT1, EXT2, EXTL1, EXTL2, and EXTL3. Studies have shown that EXT gene family members have the activities of heparan sulfate-synthesizing glycosyltransferases. EXT1 and EXT2, which have been identified as causal genes for hereditary multiple exostoses, have HS-GlcAT-II and GlcNAcT-II activities. EXTL1 has GlcNAcT-II activities. activity and EXTL3 has GlcNAcT-I and -II EXTL2 has GlcNAcT-I and Nacetylgalactosaminyltransferase activities, and transfers a GlcNAc residue to the tetrasaccharide linkage region when this region is phosphorylated by a xylose kinase 1 (FAM20B) and thereby terminate chain elongation. In mice, lack of EXTL2 causes glycosaminoglycan (GAG) overproduction and structural changes of GAGs associated with pathological processes.