



Recombinant Protein Technical Manual
Recombinant Human P63/TP63/Tumor protein p63
Protein (His & GST Tag)
RPES0596

Product Data:

Product SKU: RPES0596

Size: 20µg

Species: Human

Expression host: Baculovirus-Insect Cells

Uniprot: Q9H3D4

Protein Information:

Molecular Mass: 105 kDa

AP Molecular Mass: 125 kDa

Tag: N-His & GST

Bio-activity:

Purity: > 90 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 0.3mM DTT, 20% glycerol

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: AIS;B(p51A);B(p51B);EEC3;KET;LMS;NBP;OFC8;p40;p51;p53CP;p63;p73H;p73L;RHS;SHFM4;TP53CP;TP53L;TP73L

Immunogen Information:

Sequence: Met 1-Glu 680

Background:

Tumor protein p63 is a protein also known as transformation-related protein 63, TP63, and p63. Tumor protein p63 / p63 is a member of the p53 family of transcription factors whose members P53, p63, and p73 have similar features in their gene structures and functions. An animal model, p63^{-/-} mice has been useful in defining the role p63 plays in the development and maintenance of stratified epithelial tissues. This p63 encoding protein p63 has a dramatic impact on replenishment of cutaneous epithelial stem cells and on ovarian germ cell survival. Although these two fundamental roles of p63 attest to its powerful place in development, its other functions, specifically the apparent capacity of p63, is to supervise the emergence of new cell populations in the breast, prostate, cervix, and upper reproductive tract. P63^{-/-} mice have several development defects which include the lack of limbs and other tissues, such as teeth and mammary glands, which develop as a result of interactions between mesenchyme and epithelium. Mutations in this protein are associated with ectodermal dysplasia, and cleft lip / palate syndrome 3, ADULT syndrome (acro-dermato-ungual-lacrimal-tooth), limb-mammary syndrome, et al.