



Recombinant Protein Technical Manual

Recombinant Rat EphA4 Protein (His Tag)(Active)

RPES0875

Product Data:

Product SKU: RPES0875

Size: 200µg

Species: Rat

Expression host: HEK293 Cells

Uniprot: D3ZZK3

Protein Information:

Molecular Mass: 59.8 kDa

AP Molecular Mass: 71 kDa

Tag: C-His

Bio-activity: Measured by its binding ability in a functional ELISA. Immobilized recombinant Rat EPHA4 Protein (His Tag) at 10 µg/mL (100 µl/well) can bind Recombinant rat EFNA5-Fc with a linear range of 7.82.5 ng/ml.

Purity: > 97 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: EPHA4

Immunogen Information:

Sequence: Met1-Thr547

Background:

EPH receptor A4 (ephrin type-A receptor 4), also known as EphA4, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity⁶. EphA4 is enriched on dendritic spines of pyramidal neurons in the adult mouse hippocampus, and ephrin-A3 is localized on astrocytic processes that envelop spines. Eph receptor-mediated signaling, which is triggered by ephrins⁷, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). The extracellular domain of an EphA4 interacts with ephrin ligands, which may be tethered to neighbouring cells. Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer.