

## **Recombinant Protein Technical Manual**

# Recombinant Mouse GM-CSF/CSF2 Protein (His Tag)(Active)

**RPES0920** 

Product SKU: RPES0920 Size: 10µg

**Expression host:** Human Cells **Species**: Mouse

Uniprot: P01587

**Molecular Mass:** 15.1 kDa

AP Molecular Mass: 18-30 kDa

Tag: C-6His

**Bio-activity:** Measured in a cell proliferation assay using PDC-P1 cells. The ED50 for this effect

is 4070 pg/ml.

> 95 % as determined by SDS-PAGE **Purity:** 

**Endotoxin:** < 1.0 EU per µg as determined by the LAL method.

Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Storage:

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl, pH7.4.

Reconstitution: Please refer to the printed manual for detailed information.

**Application:** Cell Culture

Granulocyte-Macrophage Colony-Stimulating Factor; GM-CSF; Colony-Stimulating Synonyms:

Factor; CSF; Molgramostin; Sargramostim; CSF2; GMCSF

### **Immunogen Information**

**Sequence:** Ala18-Lys141

### Background:

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) was initially characterized as a growth factorthat can support the in vitro colony formation of granulocyte-macrophage progenitors. It is produced by anumber of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cellsand fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophageprogenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. Onmature hematopoietic, monocytes/ macrophages and eosinophils. GM-CSF has a functional role on nonhematopoitic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF canalso stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma andadenocarcinoma cell lines.