

# Recombinant Protein Technical Manual

# Recombinant Mouse OX40/TNFRSF4 Protein (Fc Tag)(Active)

**RPES0997** 

#### **Product Data:**

**Product SKU:** RPES0997 **Size:** 10μg

Species: Mouse Expression host: Human Cells

Uniprot: P47741

#### **Protein Information:**

Molecular Mass: 48.1 kDa

AP Molecular Mass: 65-90 kDa

**Tag:** C-Fc

**Bio-activity:** Immobilized Mouse OX40L-His(Cat: PKSM041119) at 2μg/ml(100 μl/well) can bind

Mouse OX40-Fc. The ED50 of Mouse OX40-Fc is 4.204 ug/ml.

**Purity:** > 95% as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per μg as determined by the LAL method.

**Storage:** Lyophilized protein should be stored at < -20°C, though stable at room

temperature for 3 weeks. Reconstituted protein solution can be stored at  $4-7^{\circ}$ C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:** Functional ELISA

**Synonyms:** Tumor necrosis factor receptor superfamily member 4;Tnfrsf4;OX40;CD134;Txgp1

## Immunogen Information:

Sequence: Val20-Pro211

## Background:

OX40, also termed CD134 and TNFRSF4, is a T cell co-stimulatory molecule of the TNF receptor superfamily which plays a key role in the survival and homeostasis of effector and memory T cells. OX40 is expressed on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. The interaction between OX40 and OX40 ligand (OX40L) will occur when activated T cells bind to professional antigen-presenting cells (APCs). The T-cell functions, including cytokine production, expansion, and survival, are then enhanced by the OX40 costimulatory signals. OX40 signals are critical for controlling the function and differentiation of Foxp3+ regulatory T cells. OX40-OX40L interaction regulates T-cell tolerance, peripheral T-cell homeostasis, and T-cell-mediated inflammatory diseases.