

Recombinant Protein Technical Manual Recombinant Human PACSIN2/Syndapin-2 Protein (His Tag) RPES1076

Product Data:

Product SKU: RPES1076

Species: Human

Size: 10µg

Expression host: Human Cells

Uniprot: Q9UNF0

| Protain | Inform | nation |
|---------|--------|--------|
| IUUUUII | | |

| Molecular Mass: | 56.7 kDa |
|--------------------|--|
| AP Molecular Mass: | 88 kDa |
| Tag: | C-6His |
| Bio-activity: | |
| Purity: | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin: | < 1.0 EU per μg as determined by the LAL method. |
| Storage: | Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping: | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation: | Lyophilized from a 0.2 μ m filtered solution of 20mM PB,150mM NaCl,pH7.4. |
| Reconstitution: | Please refer to the printed manual for detailed information. |
| Application: | |
| Synonyms: | Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2; PACSIN2 |

Sequence: Met 1-Gln486

Background:

Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2 (PACSIN2) is a member of the PACSIN family. PACSIN2 is localized to the plasma membrane via its coiled-coil domain. PACSIN2 is widely expressed and contains one FCH domain and one SH3 domain. PACSIN2 forms homo- and hetero-aggregates with other PACSINs. PACSIN2 may play a role in vesicle formation and transport. In addition, PACSIN2 is involved in linking the actin cytoskeleton with vesicle formation by regulating tubulin polymerization.