



Recombinant Protein Technical Manual
Recombinant Human CD89/FCAR Protein (His Tag)
RPES1349

Product Data:

Product SKU: RPES1349

Size: 10µg

Species: Human

Expression host: Human Cells

Uniprot: P24071

Protein Information:

Molecular Mass: 24.5 kDa

AP Molecular Mass: 40 kDa

Tag: C-6His

Bio-activity:

Purity: > 95 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Immunoglobulin Alpha Fc Receptor; IgA Fc Receptor; CD89; FCAR;CTB-61M7.2;FcalphaRI;FCAR;XXbac-BPG230H20.5

Immunogen Information:

Sequence: Gln22-Asn227

Background:

Immunoglobulin α Fc Receptor (IgA Fc Receptor) is a member of the immunoglobulin gene superfamily. It is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens through the charged arginin residue within its transmembrane domain. IgA Fc Receptor binds both IgA1 and IgA2 with similar affinity. The site of interaction between FCAR and IgA was identified in the first extracellular domain of FCAR and the C2/C3 junction of IgA. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. FCAR is also expressed on Kupffer cells in the liver, where it was suggested to provide a second line of defense.