



Recombinant Protein Technical Manual
Recombinant Human RSK3/RPS6KA2 Protein (GST
Tag)(Active)
RPES1417

Product Data:

Product SKU: RPES1417

Size: 20µg

Species: Human

Expression host: Baculovirus-Insect Cells

Uniprot: Q15349

Protein Information:

Molecular Mass: 110 kDa

AP Molecular Mass: 110 kDa

Tag: N-GST

Bio-activity: The specific activity was determined to be 41 nmol/min/mg using synthetic RSK peptide (KRRRLSSLRA) as substrate.

Purity: > 84 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping: This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.

Formulation: Supplied as sterile 20mM Tris, 500mM NaCl, pH 7.0, 20mM GSH

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: HU-2;MAPKAPK1C;p90-RSK3;pp90RSK3;RSK;RSK3;S6K-alpha;S6K-alpha2

Immunogen Information:

Sequence: Met 1-Leu 733

Background:

Ribosomal protein S6 kinase alpha-2, also known as 90 kDa ribosomal protein S6 kinase 2, MAP kinase-activated protein kinase 1c, MAPK-activated protein kinase 1c, Ribosomal S6 kinase 3, RSK-3, RPS6KA2 and MAPKAPK1C, is a nucleus protein which belongs to the protein kinase superfamily, AGC Ser/Thr protein kinase family and S6 kinase subfamily. RPS6KA2 / RSK-3 is expressed in many tissues. Highest expression is in lung and skeletal muscle. The expression of RPS6KA2 reduced proliferation, caused G1 arrest, increased apoptosis, reduced levels of phosphorylated extracellular signal-regulated kinase and altered other cell cycle proteins. RPS6KA2 / RSK-3 contains one AGC-kinase C-terminal domain and two protein kinase domains. It forms a complex with either ERK1 or ERK2 in quiescent cells. It transiently dissociates following mitogenic stimulation. RPS6KA2 / RSK-3 is a serine/threonine kinase that may play a role in mediating the growth-factor and stress induced activation of the transcription factor CREB. RPS6KA1, RPS6KA2, RPS6KB1, RPS6KB2, and PDK1 are involved in several pathways central to the carcinogenic process, including regulation of cell growth, insulin, and inflammation.