



Recombinant Protein Technical Manual

Recombinant Rat GFRA1/GDNFRA Protein (Fc Tag)(Active)
RPES1440

Product Data:

Product SKU: RPES1440

Size: 50µg

Species: Rat

Expression host: HEK293 Cells

Uniprot: Q62997

Protein Information:

Molecular Mass: 73.8 kDa

AP Molecular Mass: 85 kDa

Tag: C-Fc

Bio-activity: Immobilized rat His-GDNF (78-211) at 10 µg/ml (100 µl/well) can bind rat GFRA1-Fc, The EC50 of rat GFRA1-Fc is 20-46.6 ng/ml.

Purity: > 90 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: GFRA1;Gdnfra;Retl1;Trnr1;GFR alpha

Immunogen Information:

Sequence: Met1-Leu445

Background:

Glial cell line derived neurotrophic factor (GDNF) Family Receptor Alpha 1 (GFRA1) is a member of the GDNF receptor family. It is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. GFRA1 is a potent survival factor for central and peripheral neurons, and is essential for the development of kidneys and the enteric nervous system. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are its binding ligand which are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. GDNF promotes the formation of a physical complex between GFRA/GDNFR α and the orphan tyrosin kinase receptor Ret, thereby inducing its tyrosine phosphorylation. The RET is a receptor tyrosine kinase representing the signal-transducing molecule of a multisubunit surface receptor complex for the GDNF, in which GFRA / GDNFR α acts as the ligand-binding component. GDNF, a distantly related member of the transforming growth factor- β (TGF- β) superfamily, and its receptor components: GFRA1, Ret and neural cell adhesion molecule (NCAM) have been recently reported to be expressed in the testis and to be involved in the proliferation regulation of immature Sertoli cells.