



Recombinant Protein Technical Manual

Recombinant Human CSF1R/CD115 Protein (aa 20-517, His Tag)(Active) RPES1503

Product Data:

Product SKU: RPES1503

Size: 10µg

Species: Human

Expression host: Human Cells

Uniprot: NP_005202.2

Protein Information:

Molecular Mass: 55.9 kDa

AP Molecular Mass: 72 kDa

Tag: C-6His

Bio-activity: Immobilized Mouse M-CSF(Cat: PKSM041111) at 10µg/ml(100 µl/well) can bind Human M-CSFR-His.

Purity: > 95 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: Macrophage colony-stimulating factor 1 receptor; CSF receptor; CSF-R; CSFR; M-CSF-R; Proto-oncogene c-Fms; CD115; CSF1R; FMS;C-FMS;FIM2;FMS;HDLS;M-CSF-R;MCSF Receptor

Immunogen Information:

Sequence: Ile20-Pro517

Background:

Macrophage colony-stimulating factor 1 receptor (CSF1R) is a member of the type III subfamily of receptor tyrosine kinases that also includes receptors for SCF and PDGF. These receptors each contain five immunoglobulin-like domains in their extracellular domain (ECD) and a split kinase domain in their intracellular region. CSF1R is expressed primarily on cells of the monocyte/macrophage lineage, dendritic cells, stem cells and in the developing placenta. CSF1 and its receptor (CSF1R, product of *c-fms* proto-oncogene) were initially implicated as essential for normal monocyte development as well as for trophoblastic implantation. It plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone and tooth development. It is required for normal male and female fertility, and for normal development of milk ducts and acinar structures in the mammary gland during pregnancy. Aberrant expression of CSF1 or CSF1R may play a role in inflammatory diseases, such as rheumatoid arthritis, glomerulonephritis, atherosclerosis, and allograft rejection.