



Recombinant Protein Technical Manual

Recombinant Human CAMK1D Protein (GST Tag)(Active)
RPES1631

Product Data:

Product SKU: RPES1631

Size: 20µg

Species: Human

Expression host: Baculovirus-Insect Cells

Uniprot: NP_705718.1

Protein Information:

Molecular Mass: 69 kDa

AP Molecular Mass: 60 kDa

Tag: N-GST

Bio-activity: The specific activity was determined to be 70 nmol/min/mg using Autocamtide-2 synthetic peptide (KKALRRQETVDAL-amide) as substrate.

Purity: > 80 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping: This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.

Formulation: Supplied as sterile 50mM Tris, 100mM NaCl, 0.5mM Reduced Glutathione, pH 8.0

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Calcium/calmodulin-dependent protein kinase type 1D; CaM kinase I delta; CaMKI-like protein kinase; CAMK1D; CaMK1 delta; CaM-K1; CKLiK

Immunogen Information:

Sequence: Met 1-Lys 385

Background:

Calcium/calmodulin-dependent protein kinase or CaM kinases are serine/threonine-specific protein kinases that are primarily regulated by the Calcium/calmodulin complex. These kinases show a memory effect on activation. CaM kinases activity can outlast the intracellular calcium transient that is needed to activate it. In neurons, this property is important for the induction of synaptic plasticity. Pharmacological inhibition of CaM kinases II blocks the induction of long-term potentiation. Upon activation, CaM kinases II phosphorylates postsynaptic glutamate receptors and changes the electrical properties of the synapse. Calcium/calmodulin-dependent protein kinase type 1D, also known as CaM kinase I delta, CaM kinase ID, CaMKI-like protein kinase, CKLiK and CAMK1D, is a member of the protein kinase superfamily and CaMK subfamily. It contains one protein kinase domain. CAMK1D is broadly expressed. It is highly and mostly expressed in polymorphonuclear leukocytes (neutrophilic and eosinophilic granulocytes) while little or no expression is observed in monocytes and lymphocytes. Engineered overexpression of CAMK1D in non-tumorigenic breast epithelial cells led to increased cell proliferation, and molecular and phenotypic alterations indicative of epithelial-mesenchymal transition (EMT), including loss of cell-cell adhesions and increased cell migration and invasion. CAMK1D is a potential therapeutic target with particular relevance to clinically unfavorable basal-like tumors.