



Recombinant Protein Technical Manual

Recombinant Mouse CDC2 Kinase/CDK1 Protein (His & GST Tag)

RPES1874

Product Data:

Product SKU: RPES1874

Size: 50µg

Species: Mouse

Expression host: Baculovirus-Insect Cells

Uniprot: P11440

Protein Information:

Molecular Mass: 61.9 kDa

AP Molecular Mass: 57 kDa

Tag: N-His-GST

Bio-activity:

Purity: > 85 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Storage: Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping: This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.

Formulation: Supplied as sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 8.0

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Cdc2;CDC2>;Cdc2a;p34<

Immunogen Information:

Sequence: Met1-Met297

Background:

CDC2, also known as CDK1, contains 1 protein kinase domain and belongs to the protein kinase superfamily, CMGC Ser/Thr protein kinase family, CDC2/CDKX subfamily. CDC2 is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with CDC2 and function as regulatory subunits. The kinase activity of CDK1 is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of CDC2 also play important regulatory roles in cell cycle control. It is required in higher cells for entry into S-phase and mitosis. CDC2 also is a cyclin-dependent kinase which displays CTD kinase activity and is required for RNA splicing. It has CTD kinase activity by hyperphosphorylating the C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit RPB1, thereby acting as a key regulator of transcription elongation. CDK1 is required for RNA splicing, possibly by phosphorylating SRSF1/SF2. It is involved in regulation of MAP kinase activity, possibly leading to affect the response to estrogen inhibitors.