



Recombinant Protein Technical Manual

Recombinant Human ICOS/AILIM Protein (Fc Tag)

RPES2009

Product Data:

Product SKU: RPES2009

Size: 10µg

Species: Human

Expression host: Human Cells

Uniprot: Q9Y6W8

Protein Information:

Molecular Mass: 40.8 kDa

AP Molecular Mass: 47 kDa

Tag: C-Fc

Bio-activity:

Purity: > 95% as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 .

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Inducible T-cell costimulator; activation-inducible lymphocyte immunomediatory molecule; CD278; AILIM; CVID1; ICOS;

Immunogen Information:

Sequence: Glu21-Phe141

Background:

Inducible T-cell costimulator, also known as activation-inducible lymphocyte immunomediatory molecule, CD278, AILIM, CVID1 and ICOS, belongs to the CD28 and CTLA4 cell surface receptor family. . ICOS contains one Ig-like V-type domain and exists as a homodimer with disulfide-linked. ICOS is highly expressed on tonsillar T-cells and can be induced by PMA and ionomycin, ICOS plays an important role in cell-cell signaling, immune responses, and regulation of cell proliferation. Defects in ICOS are the cause of immunodeficiency common variable type 1, which is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen