



Recombinant Protein Technical Manual  
Recombinant Human CEACAM5 Protein (His Tag)  
RPES2018

### Product Data:

**Product SKU:** RPES2018

**Size:** 50µg

**Species:** Human

**Expression host:** HEK293 Cells

**Uniprot:** NP\_004354.2

### Protein Information:

**Molecular Mass:** 72.8 kDa

**AP Molecular Mass:** 10010 kDa

**Tag:** C-His

**Bio-activity:**

**Purity:** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per µg as determined by the LAL method.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from sterile PBS, pH 7.4

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:**

**Synonyms:** CD66e;CEA

## Immunogen Information:

**Sequence:** Met 1-Ala 685

## Background:

CEACAM5, also known as CEA or D66e, belongs to the large CEACAM subfamily of immunoglobulin superfamily. CEACAM5 is expressed primarily by epithelial cells, and is synthesized as a glycoprotein with a MW of 180 kDa comprising 60% carbohydrate. CEACAM5 contains one Ig-like V-type domain at the N-terminus, followed by six Ig-like C2-type domain and a GPI anchor, and exists as a homodimer. CEACAM5 and CEACAM6 are overexpressed in many cancers and are associated with adhesion and invasion. CEACAM5 can mediate cell-cell adhesion through homotypic and heterotypic interactions. It functions as a homotypic intercellular adhesion molecule and serves as a widely used tumor marker, since it is expressed at higher levels in tumorous tissues than in corresponding normal tissues. CEACAM5 has also been shown to contribute to tumorigenicity by inhibiting cellular differentiation. In addition, CEACAM5 is identified as the host receptor for the Dr family of adhesins of E. Coli, and the binding of E. coli Dr adhesins leads to dissociation of the CEACAM5 homodimer.