

Recombinant Protein Technical Manual Recombinant Mouse Vitronectin/VTN Protein (His Tag)(Active)

Product Data:

Product SKU: RPES2212

Species: Mouse

**Size:** 50µg

**RPES2212** 

Expression host: HEK293 Cells

**Uniprot:** NP\_035837.1

Protein	Inform	ation
Protein		

Molecular Mass:	54.2 kDa
AP Molecular Mass:	75-85 kDa
Tag:	C-His
Bio-activity:	Measured by the ability of the immobilized protein to support the adhesion of DU145 human prostate carcinoma cells. When cells are added to mouse Vitronectin coated plates (10 $\mu$ g/mL and 100 $\mu$ L/well), > 60% cells will adhere specifically after 30 minutes at
Purity:	> 94 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU per $\mu g$ of the protein as determined by the LAL method.
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	AI256434;Vn

## **Immunogen Information:**

## Sequence: Met 1-Lys 478

## Background:

Vitronectin, also known as VTN, is a member of the pexin family. It is an abundant glycoprotein found in serum the extracellular matrix and promotes cell adhesion and spreading. Vitronectin is a secreted protein and exists in either a single chain form or a cleaved, two chain form held together by a disulfide bond. Vitronectin is a plasma glycoprotein implicated as a regulator of diverse physiological process, including blood coagulation, fibrinolysis, pericellular proteolysis, complement dependent immune responses, and cell attachment and spreading. Because of its ability to bind platelet glycoproteins and mediate platelet adhesion and aggregation at sites of vascular injury, vitronectin has become an important mediator in the pathogenesis of coronary atherosclerosis. As a multifunctional protein with a multiple binding domain, Vitronectin interacts with a variety of plasma and cell proteins. Vitronectin binds multiple ligands, including the soluble vitronectin receptor. It may be an independent predictor of adverse cardiovascular outcomes following acute stenting. Accordingly, Vitronectin is suggested to be involved in hemostasis, cell migration, as well as tumor malignancy.